

Wave equations on space-times of low regularity: Existence results and regularity theory in the framework of generalized function algebras

James D.E. Grant & Eberhard Mayerhofer*

November 13, 2007

Abstract

We present recent developments concerning Lorentzian geometry in algebras of generalized functions. These have, in particular, raised a new interest in refined regularity theory for the wave equation on singular space-times.

Keywords: wave equation, generalised Lorentzian geometry, algebras of generalized functions.

MSC 2000: 35L05, 58J45.

1 Introduction

Generalized function algebras have proved themselves a valuable tool for non-linear analysis of partial differential equations while maintaining consistency with Schwartz' distributional framework ([9, 18] and the references therein). Their essential feature is they admit the space of smooth functions as a faithful subalgebra, which, in view of the Schwartz impossibility result [19], is the optimal possibility.

More recent scientific work also focuses on developing a geometric theory of generalized functions ([5, 12] et al.). From the very beginning of this line of research, one had in mind applications to fields such as general relativity. The foundations of generalised pseudo-Riemannian geometry were laid in [14], with further work investigating flows of generalized vector fields [13], and generalised connections on principal bundles [15].

The work of Vickers and Wilson [20] on the wave-equation on conical space-times has initiated research on existence and uniqueness results for the initial value problem of the wave-equation for a wider class of singular space-times. A first generalization of their result concerned static-space times [16] and, just recently, a quite general existence and uniqueness result has been established, thus dropping the condition of staticity [8]. In the course of this work, it became imperative to study generalized Lorentzian geometry [17], cf. below.

The program of this article is to report on recent developments in Lorentzian geometry in a generalised framework and to present results on the wave-equation on singular space-times. Finally, we suggest that regularity information for the wave-equation can be retrieved from refined energy estimates. This study is motivated by concrete examples, cf. [7].

For notation in the field, we refer to the standard reference [9].

*Faculty of Mathematics, University of Vienna, Nordbergstrasse 15, 1090 Vienna, Austria, email: {james.grant, eberhard.mayerhofer}@univie.ac.at; supported by FWF research grants P16742-N04 and Y237-N13.

2 Lorentzian geometry in algebras of generalized functions

Loosely speaking, a singular pseudo-Riemannian metric is modelled as a net of smooth metrics,

$$(g_\varepsilon)_\varepsilon \mod \mathcal{N}(\mathcal{T}_2^0(M)),$$

where g_ε is a pseudo-Riemannian metric of fixed index for sufficiently small smoothing parameter ε . As generalized functions are not determined pointwise, but only by evaluation on generalized points, one can characterize candidates for generalized pseudo-Riemannian metrics in terms of bilinear forms on the finite dimensional module $\widetilde{\mathbb{R}}^n$ (for more details, see [14], Proposition 2. 1 and [17]).

Theorem 2.1. *Let $g \in \mathcal{G}_2^0(X)$. The following are equivalent:*

1. *For each chart (V_α, ψ_α) and each $\tilde{x} \in (\psi_\alpha(V_\alpha))_\mathbb{C}^\sim$ the map $g_\alpha(\tilde{x}) : \widetilde{\mathbb{R}}^n \times \widetilde{\mathbb{R}}^n \rightarrow \widetilde{\mathbb{R}}$ is symmetric and non-degenerate.*
2. *$g : \mathcal{G}_1^0(X) \times \mathcal{G}_1^0(X) \rightarrow \mathcal{G}(X)$ is symmetric and $\det(g)$ is invertible in $\mathcal{G}((\psi_\alpha(V_\alpha)))$.*
3. *$\det g$ is invertible in $\mathcal{G}((\psi_\alpha(V_\alpha)))$ and for each relatively compact open set $V \subset X$ there exists a representative $(g_\varepsilon)_\varepsilon$ of g and $\varepsilon_0 > 0$ such that $g_\varepsilon|_V$ is a smooth pseudo-Riemannian metric for all $\varepsilon < \varepsilon_0$.*

A generalized pseudo Riemannian metric then is defined as a non-degenerate symmetric $g \in \mathcal{G}_2^0(X)$ with locally constant index. We were motivated by item (2) to continue the research in the direction of elaborating a concept of causality from the viewpoint of bilinear forms on $\widetilde{\mathbb{R}}^n$ on the basis of positivity concepts revisited in the introduction. The first step is, based on a basic perturbation theory, to introduce a well defined index associated to a symmetric matrix $A \in M(n \times n, \widetilde{\mathbb{R}})$: First, “ordered eigenvalues” of A are defined, $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. Let $j \geq 0$ such that $\lambda_1 < 0, \dots, \lambda_j < 0, \lambda_{j+1} > 0, \dots, \lambda_n > 0$, then j is called the *index* of A . (This index need not exist.) Finally, the index of a symmetric bilinear form b is then defined as the index of $A := b(e_i, e_j)$, e_i determining the canonical basis of $\widetilde{\mathbb{R}}^n$. A symmetric bilinear form b is called

- positive definite, if its index is 0.
- Lorentzian, if its index is 1.

Causality is introduced in the following way: Let b be Lorentzian. A vector $v \in \widetilde{\mathbb{R}}^n$ is called

- time-like, if $b(v, v) < 0$,
- null, if $b(v, v) = 0$,
- space-like, if $b(v, v) > 0$

Unlike the standard theory, this concept is not a trichotomy (i.e., there are vectors which are neither time-like, null or space-like), since $<<$ does not give rise to a total ordering $<=<$. Essential in the following natural basic considerations is the characterization of free vectors:

Theorem 2.2. *Let v be an element of $\widetilde{\mathbb{R}}^n$. The following are equivalent:*

1. *For any positive definite symmetric bilinear form h on $\widetilde{\mathbb{R}}^n$ we have*

$$h(v, v) > 0$$

2. *The coefficients of v with respect to some (hence any) basis span $\widetilde{\mathbb{R}}$.*
3. *v is free.*

4. For each representative $(v_\varepsilon)_\varepsilon \in \mathcal{E}_M(\mathbb{R}^n)$ of v there exists some $\varepsilon_0 \in I$ such that for each $\varepsilon < \varepsilon_0$ we have $v_\varepsilon \neq 0$ in $\mathbb{R}^n$.

Further consequences concern, for instance, characterizations of symmetric positive forms, also, direct summands, the algebraic structure of $\widetilde{\mathbb{R}}^n$ (being not semi-simple), etc. In order to establish so-called dominant energy conditions for a class of super-energy tensors, it was necessary to establish the inverse Cauchy-Schwarz inequality for time-like vectors in a Lorentzian space-time. We refer the interested reader to the foundational paper [17].

3 The wave-equation

The results of the preceding section have been elaborated in close relation with the project of generalising the paper of Vickers and Wilson [20] for the wave equation on conical space-times. Only the new framework allows to formulate our recent result, as proved in ([8]). We give here a brief overview. The initial value problem under consideration corresponds to the scalar wave equation,

$$\begin{aligned}\square_{\mathbf{g}} u &= f \\ u_\Sigma &= v \\ \widehat{\xi} u_\Sigma &= w,\end{aligned}\tag{3.1}$$

where $\square_{\mathbf{g}}$ is the d'Alembertian induced by a generalized Lorentzian metric $\mathbf{g}$ on a smooth manifold M . Here ξ is a C^∞ vector field, and $\widehat{\xi}$ the corresponding unit field; further, Σ is the space-like (in the generalized sense) initial surface. The main theorem in our joint paper with R. Steinbauer [8] establishes local existence of scalar solutions to (3.1), under the following regularity assumptions (cf. also the conclusion section of that paper)

1. For all K compact in M , for all orders of derivative $k \in \mathbb{N}_0$ and all k -tuples of vector fields $\eta_1, \dots, \eta_k \in \mathfrak{X}(M)$ and for any representative $(\mathbf{g}_\varepsilon)_\varepsilon$ we have:
 - $\sup_{p \in K} \|\mathcal{L}_{\eta_1} \dots \mathcal{L}_{\eta_k} \mathbf{g}_\varepsilon\|_m = O(\varepsilon^{-k}) \quad (\varepsilon \rightarrow 0);$
 - $\sup_{p \in K} \|\mathcal{L}_{\eta_1} \dots \mathcal{L}_{\eta_k} \mathbf{g}_\varepsilon^{-1}\|_m = O(\varepsilon^{-k}) \quad (\varepsilon \rightarrow 0).$
2. For all K compact in M , we have

$$\sup_{p \in K} \|\nabla^\varepsilon \xi_\varepsilon\|_{\mathbf{m}} = O(\log(\varepsilon)), \quad (\varepsilon \rightarrow 0),$$

where ∇^ε denotes the covariant derivative with respect to the Lorentzian metric $\mathbf{g}_\varepsilon$.

3. a causality assumption, to guarantee smooth solutions.

Different arguments suggest that the regularity estimates derived in [8] might be improved substantially by refined energy estimates based on the adaption of suitable Sobolev embedding theorems (see, e.g., [1, 10]) to our setting. In order to apply these results, we require better control on the asymptotic growth of Sobolev-constants corresponding to the regularizing sequences of Riemannian metrics induced on $t = \text{constant}$ slices of our Lorentzian space-time. This, in turn, requires control on the injectivity radius of the induced Riemannian metrics, which is possible if we have control on the diameter, curvature and volume [4]. Investigations on this topic are underway.

3.1 Regularity issues 1: Wave equations for Lorentzian metrics of Hölder regularity

The class of metrics in the preceding section can be considered as generalized Hölder-Zygmund regular of order $s = 0$ (cf. [11] and, for an adaption to the geometric setting, cf. [7]). In the

latter reference, we have investigated a two dimensional space-time $(\mathbb{R}^2, \mathbf{g})$, with $\mathbf{g}$ a Lorentzian metric of classic Hölder regularity $C^{0,\alpha}$, $\forall \alpha > 0$. The latter is designed in such a way that the principal symbol of its induced d'Alembertian $\square_{\mathbf{g}}$ is precisely the strictly hyperbolic operator used by Colombini & Spagnolo [6] (who present an example of a wave equation with Hölder continuous coefficients and smooth data, not admitting distributional solutions).

In [7] we bring the metric, by means of a coordinate transformation Φ of low regularity $C^{1,\alpha}$ ($\alpha < 1$), into a form that is evidently conformally flat. This suggests that solutions $\bar{u}$ for the wave-equation in flat space-time can then be pulled back via Φ to prospective solutions u of $\square_{\mathbf{g}} u = 0$. This procedure, however, raises questions concerning the solution concept, since the coordinate transform is not C^2 , the minimal regularity required for the standard analysis to hold. Nevertheless, for regularisations (that is, C^∞ mappings $(\Phi_\varepsilon)_\varepsilon$) of Φ , the above is rigorous and we can show that the smooth pulled back solutions $(u_\varepsilon)_\varepsilon$ even give rise to generalized solutions, that is, elements of the special algebra $\mathcal{G}$. So we have, by means of a geometric argument, found solutions of a non-trivial strictly hyperbolic differential equation with coefficients of regularity beyond Lipschitz.

3.2 Regularity issues 2: Collapsing Riemannian manifolds

An alternative direction for research concerning wave equations on manifolds with singular metrics involves situations where the induced Riemannian metric on three-dimensional hypersurfaces of the Lorentzian manifold “collapses” as $\varepsilon \rightarrow 0$. In particular, in the case of *Cheeger-Gromov collapse* [2, 3], one constructs nets of Riemannian metrics, $\mathbf{h}_\varepsilon$, with the property that, as $\varepsilon \rightarrow 0$, the sectional curvature of $\mathbf{h}_\varepsilon$ is uniformly bounded, but the injectivity radius of $\mathbf{h}_\varepsilon$ converges uniformly to zero. In this case, the volume of the three-manifold converges to zero as $\varepsilon \rightarrow 0$, but the curvature remains bounded. Taking, for example, a four-dimensional Lorentzian metric of the form $\mathbf{g}_\varepsilon = -dt^2 + \mathbf{h}_\varepsilon$, we would have asymptotic conditions

$$\left. \begin{array}{l} d\text{vol}_{\mathbf{g}_\varepsilon} = O(\varepsilon), \\ \text{Curvature} = O(1) \end{array} \right\} \quad \text{as } \varepsilon \rightarrow 0.$$

These asymptotic conditions are quite different from those investigated in [8, 20] and, as such, will lead to (generalised) solutions of the wave equation that obey quite different asymptotic conditions from those that are standard in Colombeau theory. On the other hand, Cheeger-Gromov collapse is of considerable interest from a differential geometric viewpoint, and is a field where ideas from the field of generalised functions may, perhaps, be usefully applied.

References

- [1] AUBIN, T., *Nonlinear analysis on manifolds. Monge-Ampère equations*, vol. 252 of Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, New York, 1982. 3
- [2] CHEEGER, J., GROMOV, M., *Collapsing Riemannian manifolds while keeping their curvature bounded. I*, J. Differential Geom., 23 (1986), pp. 309–346. 4
- [3] ———, *Collapsing Riemannian manifolds while keeping their curvature bounded. II*, J. Differential Geom., 32 (1990), pp. 269–298. 4
- [4] CHEEGER, J., GROMOV, M., TAYLOR, M., *Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds*, J. Differential Geom., 17 (1982), pp. 15–53. 3
- [5] COLOMBEAU, J. F., MERIL, A., *Generalized functions and multiplication of distributions on C^∞ manifolds*, J. Math. Anal. Appl., 186 (1994), pp. 357–364. 1
- [6] COLOMBINI, F. AND SPAGNOLO, S., *Some examples of hyperbolic equations without local solvability*, Ann. Sci. École Norm. Sup. (4), 22 (1989), pp. 109–125. 4
- [7] GRANT, J. D. E., MAYERHOFER, E., *Examples of wave-equations on singular space-times*, preprint. 1, 3, 4

- [8] GRANT, J. D. E., MAYERHOFER, E., STEINBAUER, R., *The wave equation on singular space-times*, preprint arXiv:0710.2007v1, (2007). [1](#), [3](#), [4](#)
- [9] GROSSER, M., KUNZINGER, M., OBERGUGGENBERGER, M., STEINBAUER, R., *Geometric Theory of Generalized Functions*, vol. 537 of Mathematics and its Applications 537, Kluwer Academic Publishers, Dordrecht, 2001. [1](#)
- [10] HEBEY, E., *Nonlinear analysis on manifolds: Sobolev spaces and inequalities*, vol. 5 of Courant Lecture Notes in Mathematics, New York University Courant Institute of Mathematical Sciences, New York, 1999. [3](#)
- [11] HÖRMANN, G., *Hölder-Zygmund regularity in algebras of generalized functions*, Z. Anal. Anwendungen, **23** (2004), pp. 139–165. [3](#)
- [12] JELÍNEK, J., *An intrinsic definition of the Colombeau generalized functions*, Comment. Math. Univ. Carolinae, **40** (1999), pp. 71–95. [1](#)
- [13] KUNZINGER, M., OBERGUGGENBERGER, M., STEINBAUER, R., VICKERS, J., *Generalized flows and singular ODEs on differentiable manifolds*, Acta Appl. Math., **80** (2004), pp. 221–241. [1](#)
- [14] KUNZINGER, M., STEINBAUER, R., *Generalized pseudo-Riemannian geometry*, Trans. Amer. Math. Soc., **354** (2002), pp. 4179–4199. [1](#), [2](#)
- [15] KUNZINGER, M., STEINBAUER, R., VICKERS, J.A., *Generalised connections and curvature*, Math. Proc. Cambridge Philos. Soc., **139** (2005), pp. 497–521. [1](#)
- [16] MAYERHOFER, E., *The wave equation on singular space-times*, PhD thesis, University of Vienna, 2006. [1](#)
- [17] ———, *On Lorentz geometry in algebras of generalized functions*, Proc. Edinb. Math. Soc., to appear, (preprint arXiv:math-ph/0604052). [1](#), [2](#), [3](#)
- [18] OBERGUGGENBERGER, M., *Multiplication of Distributions and Applications to Partial Differential Equations*, vol. **259** of Pitman Research Notes in Mathematics, Longman, Harlow, U.K., 1992. [1](#)
- [19] SCHWARTZ, L., *Sur l'impossibilité de la multiplication des distributions*, C. R. Acad. Sci. Paris, **239** (1954), pp. 847–848. [1](#)
- [20] VICKERS, J. A., WILSON, J. P., *Generalized hyperbolicity in conical spacetimes*, Class. Quantum Grav., **17** (2000), pp. 1333–1360. [1](#), [3](#), [4](#)